



DAAD

Deutscher Akademischer Austausch Dienst
German Academic Exchange Service



Ministry of Science and Education of Ukraine
National Technical University "Dnipro Polytechnic", Ukraine,
Technische Universität Bergakademie Freiberg, Germany,
under support of
GERMAN ACADEMIC EXCHANGE SERVICE (DAAD)

PROCEEDINGS
of the International Conference
"Applied Biotechnology in Mining"

April 25-27, 2018



National Technical University "Dnipro Polytechnic"
Dnipro, Ukraine

UDC 602.4:622'17:622.88(062)

Applied Biotechnology in Mining: Proceedings of the International Conference (Dnipro, April 25-27, 2018). – Dnipro: National Technical University “Dnipro Polytechnic”, – 2018. – 91 p.

ISBN

The book contains abstracts on innovative approaches for biomining technologies including both phytomining and microbial technologies in mining, as well as environmental biotechnology and other environmental assessment technologies.

The authors of the published materials are fully responsible for the selection, accuracy, reliability of the data, facts, quotations and other information.

Edited by Prof. Dr. Hermann Heilmeyer, Dr. Iryna Klimkina, Prof. Dr. Oleksandr Kovrov and others.

The Proceedings Book has been published under DAAD grant support in the frame of Program “Professionally Related Partnership with Universities of Developing Countries” and within the project “*Biotechnology in Mining – Integration of New Technologies into Educational Practice*” (2015 – 2018) due to cooperation between Technische Universität Bergakademie Freiberg (TU BAF), Germany, and National Technical University “Dnipro Polytechnic” (former National Mining University), Dnipro, Ukraine.

ISBN

© National Technical University “Dnipro Polytechnic”, Dnipro, 2018

© Litograph, 2018

<i>Svetlana Garmash and Nikita Semenov.</i> BIOUTILIZATION OF WASTE AGROINDUSTRIAL COMPLEX FOR RECEIVING BIOGAS.....	66
<i>Ilya Tkach and Artem Pavlychenko.</i> JUSTIFICATION OF MIXTURE COMPOSITION FOR RECLAMATION OF THE LANDS DISTURBED AS A RESULT OF UNDERGROUND COAL MINING.....	67
<i>Sergii Dighar, Mykhailo Yelizarov and Tetyana Kozlovska.</i> PROSPECTS FOR OBTAINING VALUABLE PRODUCTS FROM CYANOBACTERIA BIOMASS.....	68
<i>Svitlana Lysytska, Volodymyr Gerasimenko, Vasyl Kravets and Tatiana Kholodenko.</i> RESEARCH OF SECONDARY BIOMATERIALS AS POTENTIAL ENERGY SOURCE FOR MINING LABOUR.....	69
<i>Vladymyr Sytar, Vladymyr Anisimov, Natali Mitina and Svetlana Garmash.</i> BIOEDAGRADATION OF THERMOPLASTIC POLYURETHANES.....	70
<i>Polina Chirva, Iryna Pleshkova and Marina Shamray.</i> ALTERNATIVE ENERGY SOURCES: ELECTRIC CURRENT FROM LIVING PLANTS.....	71
<i>Olena Kharlamova, Volodymyr Shmandy and Tetiana Rigas.</i> BIOTECHNOLOGICAL ASPECTS USE OF WASTE IN ECOLOGICAL SAFETY.....	72
<i>Katerina Kalinina and Oleksandr Kovrov.</i> THE INFLUENCE OF MERCURY POLLUTION ON THE GROWTH OF BIO-INDICATOR PLANTS.....	73
<i>Ksenia Tsunik and Oleksandr Kovrov.</i> INVESTIGATION OF PHYTOINDICATION EFFECTS DUE TO CADMIUM CONTAMINATION.....	74
<i>Yaroslav Romanko, Irina Reshetnyak, Elena Matukhno and Angelica Meshkova.</i> INFLUENCE OF GREEN PLANTS ON DISTRIBUTION OF SOLAR RADIATION UNDER CONDITIONS OF URBAN BUILDING.....	75
<i>Kateryna Zharan and Jan C. Bongaerts.</i> THEORETICAL MODEL INTEGRATING OF RENEWABLE ENERGY INTO THE MINING INDUSTRY.....	76
Session 4. OTHER ENVIRONMENTAL TECHNOLOGIES.....	77
<i>Mykola Kharytonov and Galyna Yevgrashkyna.</i> ENVIRONMENT ASSESSMENT AND MODELLING OF SALTS TRANSFER IN FOUR LAND RESTORATION SCHEMES IN THE COAL MINING REGION.....	78
<i>Iryna Kochmar, Vasyl Karabyn and Kateryna Stepova.</i> COPPER IN COAL MINING PRODUCTS OF VIZEYSKA LVTV-VOLYN COAL BASIN MINE: ECOLOGICAL, GEOCHEMICAL AND TOXICOLOGICAL ASPECTS.....	79
<i>Tatiana Serebych and Iryna Klimkina.</i> NEUTRALIZATION EFFECT OF THE FLY ASH AND COAL DUMP WASTES.....	80
<i>Mohamed Azzaoui, Benchohra Maamar, Leila Soudani, Abassia Ayache and Mohamed Berreyah.</i> CHRONOLOGICAL STUDY OF THE VEGETATION OF FOREST MASSIF OF SDAMAS.....	81
<i>Victor Kostenko, Olena Zaviylova and Olga Chepak.</i> ECOLOGIC APPROACH TO THE RESTORATION PROCESS AT AREAS DEGRADED BY MINING OPERATIONS.....	82
<i>Nataliia Podpriatova, Iryna Klimkina and Hermann Heilmeyer.</i> STUDYING THE POSSIBILITY OF USING RED-BROWN CLAY FOR RECLAMATION OF COAL WASTE DUMPS.....	83
<i>Uliana Khromyak and Andriy Tarnavskiy.</i> REPEATING POLYETHYLENE DERIVATIVES OF THE LOW DENSE OF HRYBVOVITSKY DUMP.....	84

COPPER IN COAL MINING PRODUCTS OF VIZEYSKA LVIV-VOLYN COAL BASIN MINE: ECOLOGICAL, GEOCHEMICAL AND TOXICOLOGICAL ASPECTS

Iryna Kochmar, Vasyl Karabyn and Kateryna Stepova

*Lviv State University of Life Safety, Lviv, Ukraine
trymalevytska1@gmail.com*

Coal producers create special natural and man-made systems, which spread over large areas and affect all components of the biosphere: living and non living, biogenic and abiogenic matters. An important aspect of such a natural and man-made system is the distribution of individual chemical elements in concentrations much higher than background values. In the ecological sense, distribution of bioavailable forms of heavy metals, that can be determined in water-soluble and acetate-ammonia extracts, is the matter of high importance.

The authors have investigated the distribution of various forms of copper within the waste heap of the Vizeyska mine (Lviv-Volyn Coal Basin). This metal belongs to the second class of danger. The metal content was separately determined in the rocks of different lithological composition and the degree of thermal transformations: burnt and unburnt samples of argillites, siltstones, sandstones and soils at a distance of 50 and 100 m from the heap. Sampling was carried out in accordance with all-Union State Standard 17.4.4.02-84, preparation of solutions of extractors from the heap rock and soil was carried out in accordance with State Standards of Ukraine 4770.9: 2007. Copper concentration in extracts was determined by the atomic absorption method using the AAC-115-M-1 spectrometer.

The concentration of copper in acetate-ammonia extract ranges from 0.5 mg/kg in burnt siltstone to 5.49 mg/kg in unburnt argillite; in soils its amount decreases from 0.28 to 0.19 mg/kg depending on the distance from the heap. In aqueous extract, the concentration of Cu varies within the following limits: in unburnt argillite 2,7 mg/kg; in burned argillite, unburnt siltstone, burned siltstone, burned sandstone and unburnt coal <0.1 mg / kg (below detection limit); in the soil at a distance of 50 and 200 m from the foot of the heap < 0.1 mg / kg (below detection limit).

The maximum allowable concentration (MAC) of copper labile forms (in acetate-ammonium buffer extract) in soils is 3.0 mg/kg. The concentration of copper in soils at a distance of 50–100 m does not exceed the corresponding MAC. However, high concentrations of copper in unburnt argillite (up to 5.49 mg/kg) poses significant risks to the environmental safety of the research area.

Despite the general tolerance of plant species and genotypes to this element, copper is considered to be highly toxic. Among other factors the indirect toxic effect of copper on winter wheat is determined by varying the intensity of Ca^{2+} -dependent metabolic processes that determine the regulation of cell division and root growth. Copper is highly toxic to most fresh invertebrates. Therefore, the MAC of drinking water (European standard – 0.05, USA and CIS – 1.0, WHO – 2.0 mg/l) is higher than the fishing MAC – 0.01 mg/l. Using modified Allium test A. Dovgalyuk found that the effective toxic concentrations of copper are 23 times lower than the approved MAC of this metal in the soil and is 0.13 mg/liter.

In the human body prolonged absorption of copper along with disturbances of metabolic processes may cause cirrhosis of the liver. There are some reports about the influence of copper on the metabolism of artificially fed infants. Acute poisoning of people in the case of consuming copper with drinking water in doses of 0.14 mg/kg and above was reported.

Conclusions. Copper concentration in the acetate-ammonia extract from the heap rocks of the Vizeyska mine (Lviv-Volyn Coal Basin) varies from 0.5 mg/kg to 5.49 mg/kg and from 0.28 to 0.19 mg/kg in acetate-ammonia extract from soils in the mine area. That may be dangerous for the state of the environment.

Key words: Lviv-Volyn Coal Basin, Copper, Waste Heap of the Vizeyska Mine, Bioavailability