

Lecture Notes on Data Engineering  
and Communications Technologies 77

Sergii Babichev  
Volodymyr Lytvynenko *Editors*



# Lecture Notes in Computational Intelligence and Decision Making

2021 International Scientific  
Conference "Intellectual Systems  
of Decision-making and Problems  
of Computational Intelligence",  
Proceedings

# **Lecture Notes on Data Engineering and Communications Technologies**

Volume 77

## **Series Editor**

Fatos Xhafa, Technical University of Catalonia, Barcelona, Spain

The aim of the book series is to present cutting edge engineering approaches to data technologies and communications. It will publish latest advances on the engineering task of building and deploying distributed, scalable and reliable data infrastructures and communication systems.

The series will have a prominent applied focus on data technologies and communications with aim to promote the bridging from fundamental research on data science and networking to data engineering and communications that lead to industry products, business knowledge and standardisation.

Indexed by SCOPUS, INSPEC, EI Compendex.

All books published in the series are submitted for consideration in Web of Science.

More information about this series at <http://www.springer.com/series/15362>


Sergii Babichev · Volodymyr Lytvynenko  
Editors

# Lecture Notes in Computational Intelligence and Decision Making

2021 International Scientific Conference  
“Intellectual Systems of Decision-making  
and Problems of Computational Intelligence”,  
Proceedings

 Springer

*Editors*

Sergii Babichev   
Department of Physics  
Kherson State University  
Kherson, Ukraine

Volodymyr Lytvynenko  
Department of Informatics  
and Computer Science  
Kherson National Technical University  
Kherson, Ukraine

ISSN 2367-4512                      ISSN 2367-4520 (electronic)  
Lecture Notes on Data Engineering and Communications Technologies  
ISBN 978-3-030-82013-8              ISBN 978-3-030-82014-5 (eBook)  
<https://doi.org/10.1007/978-3-030-82014-5>

© Springer Nature Switzerland AG 2022

This work is subject to copyright. All rights are solely and exclusively licensed by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, expressed or implied, with respect to the material contained herein or for any errors or omissions that may have been made. The publisher remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This Springer imprint is published by the registered company Springer Nature Switzerland AG  
The registered company address is: Gewerbestrasse 11, 6330 Cham, Switzerland

# Contents

<b>Analysis and Modeling of Complex Systems and Processes</b>	
<b>Financial Risk Estimation in Conditions of Stochastic Uncertainties . . .</b>	<b>3</b>
Oleksandr Trofymchuk, Peter Bidyuk, Irina Kalinina, and Aleksandr Gozhyj	
<b>Numerical Modeling of Disk Dissolution in Melt During Gas Blowing . . . . .</b>	<b>25</b>
Kyrylo Krasnikov	
<b>Streaming Algorithm to the Decomposition of a Polyatomic Molecules Mass Spectra on the Polychlorinated Biphenyls Molecule Example . . . .</b>	<b>39</b>
Serge Olszewski, Violetta Demchenko, Eva Zaets, Volodymyr Lytvynenko, Irina Lurie, Oleg Boskin, and Sergiy Gnatyuk	
<b>Method of Functional-Value Calculations of Complex Systems with Mixed Subsystems Connections . . . . .</b>	<b>54</b>
Maksym Korobchynskyi, Mykhailo Slonov, Pavlo Krysiak, Myhailo Rudenko, and Oleksandr Maryliv	
<b>Current State of Methods, Models, and Information Technologies of Genes Expression Profiling Extraction: A Review . . . . .</b>	<b>69</b>
Lyudmyla Yasinska-Damri, Ihor Liakh, Sergii Babichev, and Bohdan Durnyak	
<b>A Method of Analytical Calculation of Dynamic Characteristics of Digital Adaptive Filters with Parallel-Sequential Weight Summation . . .</b>	<b>82</b>
Kostiantyn Semibalamut, Volodymyr Moldovan, Svitlana Lysenko, Maksym Topolnytskyi, and Sergiy Zhuk	
<b>Simulation Modeling as a Means of Solving Professionally-Oriented Problems in Maritime Industry . . . . .</b>	<b>94</b>
Tatyana Zaytseva, Lyudmyla Kravtsova, Oksana Tereshchenkova, and Alona Yurzhenko	

<b>Data Mining Methods, Models and Solutions for Big Data Cases in Telecommunication Industry</b> . . . . .	107
Nataliia Kuznietsova, Peter Bidyuk, and Maryna Kuznietsova	
<b>Agile Architectural Model for Development of Time-Series Forecasting as a Service Applications</b> . . . . .	128
Illia Uzun, Ivan Lobachev, Luke Gall, and Vyacheslav Kharchenko	
<b>Essential <math>R</math> Peak Detector Based on the Polynomial Fitting</b> . . . . .	148
Olga Velychko, Oleh Datsok, and Iryna Perova	
<b>An Approach to Identifying and Filling Data Gaps in Machine Learning Procedures</b> . . . . .	164
Peter Bidyuk, Irina Kalinina, and Aleksandr Gozhyy	
<b>Information Technology for Assessing the Situation in Energy-Active Facilities by the Operator of an Automated Control System During Data Sampling</b> . . . . .	177
Lubomyr Sikora, Natalya Lysa, Roman Martysyshyn, and Yuliya Miyushkovych	
<b>Application of Ensemble Methods of Strengthening in Search of Legal Information</b> . . . . .	188
Nataliya Boyko, Khrystyna Kmetyk-Podubinska, and Iryna Andrusiak	
<b>Synthesis of Barker-Like Codes with Adaptation to Interference</b> . . . . .	201
Oleg Riznyk, Ivan Tsmots, Roman Martysyshyn, Yuliya Miyushkovych, and Yurii Kynash	
<b>Models of Factors of the Design Process of Reference and Encyclopedic Book Editions</b> . . . . .	217
Vsevolod Senkivskyy, Iryna Pikh, Alona Kudriashova, Nataliia Senkivska, and Lyubov Tupyachak	
<b>Analysis of Digital Processing of the Acoustic Emission Diagnostics Informative Parameters Under Deformation Impact Conditions</b> . . . . .	230
Volodymyr Marasanov, Hanna Rudakova, Dmitry Stepanchikov, Oleksandr Sharko, Artem Sharko, and Tetiana Kiryushatova	
<b>Solution of the Problem of Optimizing Route with Using the Risk Criterion</b> . . . . .	252
Pavlo Mamenko, Serhii Zinchenko, Vitaliy Kobets, Pavlo Nosov, and Ihor Popovych	
<b>Automatic Optimal Control of a Vessel with Redundant Structure of Executive Devices</b> . . . . .	266
Serhii Zinchenko, Oleh Tovstokoryi, Andrii Ben, Pavlo Nosov, Ihor Popovych, and Yaroslav Nahrybelnyi	

**Practice Analysis of Effectiveness Components for the System Functioning Process: Energy Aspect** . . . . . 282  
 Victor Yarmolenko, Nataliia Burennikova, Sergii Pavlov, Vyacheslav Kavetskiy, Igor Zavgorodnii, Kostiantyn Havrysh, and Olga Pinaieva

**Method of Mathematical and Geoinformation Models Integration Based On Unification of the Ecological Data Formalization** . . . . . 297  
 Oleg Mashkov, Taras Ivashchenko, Waldemar Wójcik, Yuriy Bardachov, and Viktor Kozel

**Prediction of Native Protein Conformation by a Hybrid Algorithm of Clonal Selection and Differential Evolution** . . . . . 314  
 Iryna Fefelova, Andrey Fefelov, Volodymyr Lytvynenko, Oksana Ohnieva, and Saule Smailova

**Reduction of Training Samples in Solar Insolation Prediction Under Weather and Climatic Changes** . . . . . 331  
 Yakiv Povod, Volodymyr Sherstjuk, and Maryna Zharikova

**Research of Acoustic Signals Digital Processing Methods Application Efficiency for the Electromechanical System Functional Diagnostics** . . . 349  
 Hanna Rudakova, Oksana Polyvoda, Inna Kondratieva, Vladyslav Polyvoda, Antonina Rudakova, and Yuriy Rozov

**Computer Simulation of Physical Processes Using Euler-Cromer Method** . . . . . 367  
 Tatiana Goncharenko, Yuri Ivashina, and Nataliya Golovko

**Automated System and Domain-Specific Language for Medical Data Collection and Processing** . . . . . 377  
 Oleksii Boiarskyi and Svitlana Popereshnyak

**Applying Visibility Graphs to Classify Time Series** . . . . . 397  
 Lyudmyla Kirichenko, Tamara Radivilova, and Vitalii Ryzhanov

**Theoretical and Applied Aspects of Decision-Making Systems**

**Assessment of the Influencing Factors Significance in Non-destructive Testing Systems of Metals Mechanical Characteristics Based on the Bayesian Network** . . . . . 413  
 Volodymyr Mirnenko, Oleksandr Mishkov, Anatolii Balanda, Vasily Nadraga, and Oleksandr Hryhorenko

**Markovian Learning Methods in Decision-Making Systems** . . . . . 423  
 Petro Kravets, Yevhen Burov, Vasyi Lytvyn, Victoria Vysotska, Yuriy Ryshkovets, Oksana Brodyak, and Svitlana Vyshemyrska



**Fine-Tuning of the Measure for Full Reference Image Quality Assessment** . . . . . 438  
 Oleksii Gorokhovatskyi and Olena Peredrii

**A Model for Assessing the Rating of Higher Education School Academic Staff Members Based on the Fuzzy Inference System** . . . . . 449  
 Sergii Babichev, Aleksander Spivakovsky, Serhii Omelchuk, and Vitaliy Kobets

**Early Revealing of Professional Burnout Predictors in Emergency Care Workers** . . . . . 464  
 Igor Zavgorodnii, Olha Lalymenko, Iryna Perova, Polina Zhernova, Anastasiia Kiriak, and Oleksandr Novytsky

**Forming Predictive Features of Tweets for Decision-Making Support** . . . . . 479  
 Bohdan M. Pavlyshenko

**Method for Adaptive Semantic Testing of Educational Materials Level of Knowledge** . . . . . 491  
 Olexander Mazurets, Olexander Barmak, Iurii Krak, Eduard Manziuk, and Ruslan Bahrii

**Baseline Wander Correction of the Electrocardiogram Signals for Effective Preprocessing** . . . . . 507  
 Anatolii Pashko, Iurii Krak, Oleg Stelia, and Waldemar Wojcik

**Intellectual Information Technologies of the Resources Management in Conditions of Unstable External Environment** . . . . . 519  
 Marharyta Sharko, Olga Gonchar, Mykola Tkach, Anatolii Polishchuk, Nataliia Vasylenko, Mikhailo Mosin, and Natalia Petrushenko

**Information Technologies and Neural Network Means for Building the Complex Goal Program “Improving the Management of Intellectual Capital”** . . . . . 534  
 Anzhelika Azarova

**Quantitative Assessment of Forest Disturbance with C-Band SAR Data for Decision Making Support in Forest Management** . . . . . 548  
 Anna Kozlova, Sergey Stankevich, Mykhailo Svideniuk, and Artem Andreiev

**An Intelligent System for Providing Recommendations on the Web Development Learning** . . . . . 563  
 Iryna Yurchuk and Mykyta Kutsenko

**Real-Time Sensing, Reasoning and Adaptation for Computer Vision Systems** . . . . . 573  
 Volodymyr Hrytsyk and Mariia Nazarkevych

**Computational Intelligence and Inductive Modeling**

**Spectrophotometric Method for Coagulant Determining in a Stream Based on an Artificial Neural Network** ..... 589  
 Andrii Safonyk, Maksym Mishchanchuk, and Ivanna Hrytsiuk

**Comparative Analysis of Normalizing Techniques Based on the Use of Classification Quality Criteria** ..... 602  
 Oleksandr Mishkov, Kostiantyn Zorin, Denys Kovtoniuk, Vladyslav Dereko, and Igor Morgun

**Robust Recurrent Credibilistic Modification of the Gustafson - Kessel Algorithm** ..... 613  
 Yevgeniy Bodyanskiy, Alina Shafronenko, Iryna Klymova, and Vladyslav Polyvoda

**Tunable Activation Functions for Deep Neural Networks** ..... 624  
 Bohdan Bilonoh, Yevgeniy Bodyanskiy, Bohdan Kolchygin, and Sergii Mashtalir

**Markov-Chain-Based Agents for k-Armed Bandit Problem** ..... 634  
 Vladyslav Sarnatskyi and Igor Baklan

**Predicting Customer Churn Using Machine Learning in IT Startups** ..... 645  
 Viktor Morozov, Olga Mezentseva, Anna Kolomiiets, and Maksym Proskurin

**Union of Fuzzy Homogeneous Classes of Objects** ..... 665  
 Dmytro Terletskyi and Sergey Yershov

**Neuro-Fuzzy Diagnostics Systems Based on SGTm Neural-Like Structure and T-Controller** ..... 685  
 Roman Tkachenko, Ivan Izonin, and Pavlo Tkachenko

**An Integral Software Solution of the SGTm Neural-Like Structures Implementation for Solving Different Data Mining Tasks** ..... 696  
 Roman Tkachenko

**An Expert System Prototype for the Early Diagnosis of Pneumonia** ... 714  
 Mariia Voronenko, Olena Kovalchuk, Luidmyla Lytvynenko, Svitlana Vyshemyska, and Iurii Krak

**Using Bayesian Networks to Estimate the Effectiveness of Innovative Projects** ..... 729  
 Oleksandr Naumov, Mariia Voronenko, Olga Naumova, Nataliia Savina, Svitlana Vyshemyska, Vitaliy Kornychuk, and Volodymyr Lytvynenko

**Method of Transfer Deep Learning Convolutional Neural Networks for Automated Recognition Facial Expression Systems . . . . . 744**  
Arsirii Olena, Denys Petrosiuk, Babilunha Oksana,  
and Nikolenko Anatolii

**Development of a Smart Education System for Analysis and Prediction of Students' Academic Performance . . . . . 762**  
Svetlana Yaremko, Elena Kuzmina, Nataliia Savina, Dmitriy Yaremko,  
Vladyslav Kuzmin, and Oksana Adler

**Assesment Model for Domain Specific Programming Language Design . . . . . 776**  
Oleksandr Ocheretianyi and Ighor Baklan

**General Scheme of Modeling of Longitudinal Oscillations in Horizontal Rods . . . . . 789**  
Roman Tatsij, Oksana Karabyn, Oksana Chmyr, Igor Malets,  
and Olga Smotr

**Author Index . . . . . 803**



# General Scheme of Modeling of Longitudinal Oscillations in Horizontal Rods

Roman Tatsij , Oksana Karabyn <sup>(✉)</sup> , Oksana Chmyr , Igor Malets ,  
and Olga Smotr 

Lviv State University of Life Safety, Lviv, Ukraine

**Abstract.** In this paper, we present the results of modeling nonstationary oscillatory processes in rods consisting of an arbitrary number of pieces. When modeling oscillatory processes that occur in many technical objects (automotive shafts, rods) an important role is played by finding the amplitude and frequency of oscillations. Solving oscillatory problems is associated with various difficulties. Such difficulties are a consequence of the application of methods of operation calculus and methods of approximate calculations. The method of modeling of oscillatory processes offered in work is executed without application of operational methods and methods of approximate calculations. The method of oscillation process modeling proposed in this paper is a universal method. The work is based on the concept of quasi-derivatives. Applying the concept of quasi-derivatives helps to avoid the problem of multiplication of generalized functions. Analytical formulas for describing oscillatory processes in rods consisting of an arbitrary number of pieces are obtained. It can be applied in cases where pieces of rods consist of different materials, and also when in places of joints the masses are concentrated. The proposed method allows the use of computational software. An example of constructing eigenvalues and eigenfunctions for a rod consisting of two pieces is given.

**Keywords:** Kvazidifferential equation · The boundary value problem · The cauchy matrix · The eigenvalues problem · The method of fourier and the method of eigenfunctions

## 1 Introduction

The problem of finding eigenvalues and eigenfunctions for equations in partial derivatives of the second order is an urgent problem. The relevance is due to the fact that such problems arise in the modeling of oscillatory processes of many technical systems. Each specific model is a separate mathematical problem, the ability to solve which depends on the input conditions. The theory of oscillatory processes is described in detail in [2, 4, 15, 17]. In these works the mathematical and physical bases of oscillatory processes and methods of their modeling are stated. The method of solving boundary value problems according to which the solution of

problems is reduced to solving simpler problems is called the method of reduction. The solution of the general boundary value problem is sought in the form of the sum of functions, one of which is the solution of a homogeneous problem, and the other is an inhomogeneous problem with zero boundary conditions. One of the most common methods for solving inhomogeneous problems is the method of separating variables, or the Fourier method. According to this method, the solution of a homogeneous problem is sought in the form of the product of two functions of one variable. Such a problem is called a problem of eigenvalues and eigenfunctions. It is also called the task of Shturm Liouville [1, 9, 11]. The Fourier method is an accurate method of solving these problems. In the process of solving problems by this method there are problems with the justification of the convergence of series and the multiplication of generalized functions [6, 10, 19]. In some cases, these problems can be avoided by reducing the system to a matrix form by introducing the so-called quasi-derivative [4, 7, 13]. In this paper, we used this method to find solutions to four problems of oscillatory processes and demonstrated the possibility of finding the required number of eigenvalues and eigenfunctions. This method is new. It allows you to avoid multiplication of generalized functions. The proposed method has an advantage over other methods in that it allows the use of computational mathematical packages.

## 2 Problem Statement

Oscillation processes are modeled using hyperbolic type equations. Quite often, it is almost impossible to obtain closed-loop solutions of such differential equations. The method proposed in our work belongs to the direct methods of solving boundary value problems, as a result of which the solutions are obtained in a closed form. A feature of our work is the use of a quasi-derivative. This approach will make it possible to write equations with partial derivatives of the second order of the hyperbolic type with general boundary conditions to the matrix form. The process of constructing the solution is based on the multiplication of matrices. Consider a second-order differential equation in partial derivatives

$$m(x) \frac{\partial^2 u}{\partial t^2} = \frac{\partial}{\partial x} (\lambda(x) \frac{\partial u}{\partial x}) \tag{1}$$

where  $x \in (x_0; x_n)$ ,  $t \in (0; +\infty)$

Denote the product  $\lambda(x) \frac{\partial u}{\partial x}$  by  $u^{[1]}$  and let's call it quasi-derivative. Write down the general boundary conditions:

$$\begin{cases} p_{11}u(x_0; t) + p_{12}u^{[1]}(x_0; t) + q_{11}u(x_n; t) + q_{12}u^{[1]}(x_n; t) = \psi_0(t) \\ p_{21}u(x_0; t) + p_{22}u^{[1]}(x_0; t) + q_{21}u(x_n; t) + q_{22}u^{[1]}(x_n; t) = \psi_n(t) \end{cases} \tag{2}$$

The initial conditions are the following:

$$\begin{cases} u(x; 0) = \Phi_0(x) \\ \frac{\partial u}{\partial t}(x; 0) = \Phi_1(x) \end{cases} \tag{3}$$

where  $\psi_0(t), \psi_n(t) \in C^2(0; +\infty)$ ,  $\Phi_0(x), \Phi_1(x)$  are piecewise continuous on  $[x_0; x_n]$

### 3 Literature Review

As defined in the introduction, the problem of finding eigenvalues and eigenfunctions of a level in private derivatives of other hyperbolic conditions is an urgent problem in modeling quantitative processes of different technical systems. In the article [8] the case of rod oscillations under the action of periodic force is considered. What is special is that the action of force is distributed along the rod at a certain speed. Finding eigenvalues and eigenfunctions is a key task for solving the equation with second-order partial derivatives. The work [7] is devoted to finding a class of self-adjoint regular eigenfunctions and eigenvalues for each natural  $n$ . The boundary value problem for the hyperbolic equation in the rectangular domain is considered in [12]. A feature of this work is the case of singular coefficients of the equation. A very important aspect is the proof of the existence and uniqueness of the solution (Cauchy problem) and the proof of the stability theorem of the solution. The solution is obtained in the form of a Fourier-Bessel series. In the article [16] the matrix approach to the decision of a problem on eigenvalues and eigenfunctions (Sturm-Liouville problem) of the equation of hyperbolic type is offered. Emphasis is placed on the fact that the type of solution depends on the structure of the matrices. The matrix approach to the problem presentation is very convenient for the use of computational software. In our case the method of matrix calculus and the presentation of the differential equation in partial derivatives and the most general boundary conditions in matrix form are also used. In [3] a two-step method of discretization of a combined hyperbolic-parabolic problem with a nonlocal boundary condition was proposed. Examples of solving such problems by numerical methods are given. The problem of stability of solutions of the second-order problem considered in the work [18]. The method of introducing a quasi-derivative and reducing the equation of thermal conductivity to the matrix form is used in the work [13]. However, despite the achievements in this subject area, the problem of series convergence, multiplication of generalized functions and obtaining solutions of equations in partial derivatives remains open. The solution to this problem can be achieved to some extent through the use of modern calculation technologies using software tools [5]. The authors proposed to use the matrix form of the second-order differential equation and modern methods of calculations using mathematical software to model the oscillatory processes in the works [14]. The objective of the research is to find solutions of the equation of rod oscillations with different load and conditions using the concept of a quasi-derivative.

### 4 The General Scheme of Search of the Solution

The general scheme of finding a solution is to build two functions  $w(x, t)$  and  $v(x, t)$  such as

$$u(x, t) = w(x, t) + v(x, t)$$

Let's find the function  $w(x, t)$  by a constructive method, then the function  $v(x, t)$  will be defined unambiguously. The function  $w(x, t)$  is a solution of a boundary value problem

$$(\lambda(x)w_x')'_x = -f(x) \tag{4}$$

with the boundary conditions (2). We reduce this problem to a matrix form

$$\overline{W}'_x = A(x) \cdot \overline{W} + \overline{F} \tag{5}$$

$$P \cdot \overline{W}(x_0, t) + Q \cdot \overline{W}(x_n, t) = \overline{I}(t) \tag{6}$$

Where

$$A(x) = \begin{pmatrix} 0 & \frac{1}{\lambda(x)} \\ 0 & 0 \end{pmatrix}, \overline{W} = \begin{pmatrix} w \\ w^{[1]} \end{pmatrix}, \overline{F} = \begin{pmatrix} 0 \\ -f(x) \end{pmatrix},$$

$$P = \begin{pmatrix} p_{11} & p_{12} \\ p_{21} & p_{22} \end{pmatrix}, Q = \begin{pmatrix} q_{11} & q_{12} \\ q_{21} & q_{22} \end{pmatrix}, \overline{I}(t) = \begin{pmatrix} \psi_0(t) \\ \psi_1(t) \end{pmatrix}$$

Let  $x_0 < x_1 < x_2 < \dots < x_{j-1} < x_j < x_{j+1} < \dots < x_{n-1} < x_n$  are the arbitrary partition of the segment  $[x_0; x_n]$  of the real axis  $Ox$  into  $n$  parts. The solution of the problem at each interval  $[x_i; x_{i+1})$  is the following:

$$\overline{W}_i(x, t) = B_i(x, x_i) \cdot \overline{P}_i + \int_{x_i}^x B_i(x, s) \cdot \overline{P}_i(s) ds$$

where  $B_i(x, s) = \begin{pmatrix} 1 & b_i(x, s) \\ 0 & 1 \end{pmatrix}$  is the Cauchy matrix of a system,  $b_i(x, s) = \int_s^x \frac{1}{\lambda_i(z)} dz$  Let's build matrices (for an arbitrary  $k \geq i$ )

$$B(x_k, x_i) = \begin{pmatrix} 1 & \sum_{m=i}^{k-1} b_m(x_{m+1}, x_m) \\ 0 & 1 \end{pmatrix}$$

We use the recurrent method of mathematical induction to construct vectors  $\overline{P}_i$

$$\overline{P}_i = B(x_i, x_0) \cdot \overline{P}_0 + \sum_{k=0}^i B(x_i, x_k) \overline{Z}_k,$$

where  $\overline{Z}_k = \int_{x_{k-1}}^{x_k} B_{k-1}(x_k, s) \cdot \overline{F}_{k-1}(s) ds$ . The vector  $\overline{P}_0$  determines from the initial conditions by the formula:

$$\overline{P}_0 = [P + Q \cdot B(x_n, x_0)]^{-1} \cdot (\overline{I} - Q \sum_{k=1}^n B(x_n, x_k) \overline{Z}_k)$$

The first coordinate of the vector  $\overline{W}_i(x, t)$  is indeed the searched function  $w_i(x, t)$ . Thus function  $w(x, t)$  is a sum  $w(x, t) = \sum_{i=0}^{n-1} w_i(x, t) \theta_i$ ,  $\theta_i$  is the characteristic function of the interval  $[x_i; x_{i+1})$ , that is  $\theta_i(x) = \begin{cases} 1, & x \in [x_i, x_{i+1}), \\ 0, & x \notin [x_i, x_{i+1}), \end{cases} i = \overline{0, n-1}$ .

The function  $v(x, t)$  is the solution of a mixed problem

$$m(x) \frac{\partial^2 v}{\partial t^2} - \frac{\partial}{\partial x} (\lambda(x) \frac{\partial v}{\partial x}) = -m(x) \frac{\partial^2 w}{\partial t^2} \tag{7}$$

Initial conditions for the function  $v(x, t)$  are the following:

$$\begin{cases} v(x, 0) = \Phi_0(x), \\ \frac{\partial v}{\partial t}(x, 0) = \Phi_1(x), \end{cases} \tag{8}$$

where

$$\begin{aligned} \Phi_0 &\stackrel{def}{=} \phi_0(x) - w(x, 0) \\ \Phi_1 &\stackrel{def}{=} \phi_1(x) - \frac{\partial w}{\partial t}(x, 0). \end{aligned}$$

The boundary conditions for the function  $v(x, t)$  will be the following:

$$\begin{cases} p_{11}v(x_0) + p_{12}v^{[1]}(x_0) = 0 \\ q_{21}v(x_n) + q_{22}v^{[1]}(x_n) = 0 \end{cases} \tag{9}$$

We build the function  $v(x, t)$  in the form of a product

$$v(x, t) = T(t) \cdot X(x)$$

Solve the problem of eigenvalues and eigenfunctions

$$(\lambda(x)X'(x))' + \omega^2 m(x)X(x) = 0 \tag{10}$$

$$\begin{cases} X(x_0) = 0 \\ X(x_n) = 0 \end{cases} \tag{11}$$

Let's transform the problem into a matrix form. For this purposes let's denote

$$X[1] \stackrel{def}{=} \lambda X', \bar{X} = \begin{pmatrix} X \\ X[1] \end{pmatrix}, A = \begin{pmatrix} 0 & \frac{1}{\lambda(x)} \\ -m(x)\omega^2 & 0 \end{pmatrix}.$$

Cauchy matrix

$$\tilde{B}(x_n, x_0, \omega) \stackrel{def}{=} \begin{pmatrix} b_{11}(\omega) & b_{12}(\omega) \\ b_{21}(\omega) & b_{22}(\omega) \end{pmatrix}.$$

With this designations, the problem (10), (11) takes the form:

$$\bar{X}' = A \cdot X \tag{12}$$

$$P \cdot \bar{X}(x_0) + Q \cdot \bar{X}(x_n) = 0. \tag{13}$$

We solve the problem in each segment  $[x_i; x_{i+1})$ . The eigenfunctions  $X_k(x, \omega_k)$  as the first coordinates of the eigenvectors  $\bar{X}_k(x, \omega_k)$  can be written down as

$$X_k(x, \omega_k) = (1 \ 0) \cdot \tilde{B}(x, x_0, \omega_k) \cdot \bar{C} \tag{14}$$

In order to exist vector  $\bar{C}$ , there must be such condition

$$\det(P + Q \cdot \tilde{B}(x_n, x_0, \omega)) = 0 \tag{15}$$



The Eq. (14) is a characteristic equation of the problem (10), (11) or (12), (13). We will look for the function  $v(x, t)$  by arranging it in a series by orthogonal functions  $X_k(x, \omega_k)$ , which are the solutions of the set of equations

$$v(x, t) = \sum_{k=1}^n T_k(t) \cdot X_k(x, \omega_k), \tag{16}$$

where  $T_k(t)$  are the solutions of a set of equations

$$T_k''(t) + \omega_k^2 \cdot T_k(t) = -w_k(t).$$

Thus, finally a solution of the mixed problem (7)–(8) is received in a form of the series

$$v(x, t) = \sum_{k=1}^{\infty} \left[ \Phi_{0k} \cos \omega_k t + \frac{\Phi_{1k}}{\omega_k} \sin \omega_k t - \frac{1}{\omega_k} \int_0^t \sin \omega_k(t - s) \cdot w_k(s) ds \right] \cdot X_k(x, \omega_k). \tag{17}$$

For some function types consider how the method description works.

### 5 A Case of Piecewise Continuous Functions with Concentrated Masses

Let in (1) define  $m(x)$  and  $\lambda(x)$  - are piecewise continuous functions

$$m(x) = \sum_{i=0}^{n-1} m_i(x)\theta_i, \quad m_i(x) > 0,$$

$$\lambda(x) = \sum_{i=0}^{n-1} \lambda_i(x)\theta_i, \quad \lambda_i(x) \in C[x_i, x_{i+1}], \quad \lambda_i(x) > 0,$$

$$f(x) = \sum_{i=0}^{n-1} g_i(x)\theta_i + \sum_{j=0}^{n-1} s_j \delta_j(x - x_j), \quad g_i \in C[x_i, x_{i+1}], \quad s_j \in \mathbb{R},$$

$$P = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad Q = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

By using the described method, we can avoid multiplication of generalized functions that are embedded in the function  $f(x)$ . Let's denote

$$\sigma_n = \sum_{m=0}^{n-1} b_m(x_{m+1}, x_m), \quad I_{k-1}(x_k) = - \int_{x_{k-1}}^{x_k} b_{k-1}(x_k, s) g_{k-1}(s) ds,$$

$$I_{k-1}^{[1]}(x_k) = - \int_{x_{k-1}}^{x_k} g_{k-1}(s) ds.$$

With such designations

$$\begin{aligned}
 w_i(x, t) = & \psi_0(t) + (b_i(x, x_i) + \sigma_i) \cdot \frac{\psi_n(t) - \psi_0(t)}{\sigma_n} \\
 & - \frac{1}{\sigma_n} (b_i(x, x_i) + \sigma_i) \cdot \sum_{k=0}^n (I_{k-1}(x_k) + (I_{k-1}^{[1]}(x_k) - s_k) \cdot \sum_{m=k}^{n-1} b_m(x_{m+1}, x_m) \\
 & + \sum_{k=0}^i (I_{k-1}(x_k) + (I_{k-1}^{[1]}(x_k) - s_k) \sum_{m=k}^{i-1} b_m(x_{m+1}, x_m)) \\
 & + b_i(x, x_i) \sum_{k=0}^i (I_{k-1}^{[1]}(x_k) - s_k) + I_i(x)
 \end{aligned}$$

Vector  $\bar{C}$  in (14) is  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ . The characteristic equation of function  $v(x, t)$  takes the form  $b_{12}(\omega) = 0$  and function  $v(x, t)$  takes the form (17).

### 6 A Case of Piecewise Constant Functions with Concentrated Masses

Let in (1)  $m(x), f(x)$  – are piecewise constant with concentrated masses,  $\lambda(x)$  – is piecewise constant:

$$\begin{aligned}
 m(x) = & \sum_{i=0}^{n-1} m_i \theta_i(x) + \sum_{i=0}^{n-1} M_i \delta(x - x_i), \\
 f(x) = & \sum_{i=0}^{n-1} g_i \theta_i(x) + \sum_{i=0}^{n-1} S_i \delta(x - x_i),
 \end{aligned}$$

and  $\lambda(x)$  – are piecewise constant functions  $\lambda(x) = \sum_{i=0}^{n-1} \lambda_i \theta_i(x)$ . In boundary conditions (2)

$$P = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, Q = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

In this case it is possible to specify the function  $w_i(x, t)$

$$\begin{aligned}
 w_i(x, t) = & \psi_0(t) + \left( \frac{x - x_i}{\lambda_i} + \sum_{m=0}^{i-1} \frac{\Delta x_{m+1}}{\lambda_m} \right) \cdot \frac{\psi_n(t) - \psi_0(t)}{\sum_{m=0}^{n-1} \frac{\Delta x_{m+1}}{\lambda_m}} \\
 & - \frac{1}{\sum_{m=0}^{n-1} \frac{\Delta x_{m+1}}{\lambda_m}} \cdot \left( \frac{x - x_i}{\lambda_i} + \sum_{m=0}^{i-1} \frac{\Delta x_{m+1}}{\lambda_m} \right).
 \end{aligned}$$

$$\begin{aligned} & \cdot \left( \sum_{k=1}^n \frac{g_{k-1} \Delta x_k^2}{2\lambda_{k-1}} - (g_{k-1} \Delta x_k + s_k) \sum_{m=k}^{n-1} \frac{\Delta x_{m-1}}{\lambda_m} \right) \\ & + \sum_{k=1}^i \left( \frac{g_{k-1} \Delta x_k^2}{2\lambda_{k-1}} - (g_{k-1} \Delta x_k + s_k) \sum_{m=k}^{i-1} \frac{\Delta x_{m-1}}{\lambda_m} \right) \\ & - \frac{x - x_i}{\lambda_i} \sum_{k=1}^i (g_{k-1} \Delta x_k + s_k) + \frac{g_i (x - x_i)^2}{2\lambda_i} \end{aligned}$$

To build a function  $v(x, t)$  we use the method of expansion by the eigenfunctions. Due to the delta functions in the left and right part of the equation, we go to the system

$$\bar{X}' = \left( \sum_{k=0}^{n-1} A_k \theta_k + \sum_{k=0}^{n-1} C_k \delta(x - x_k) \right) \cdot \bar{X},$$

where  $A_k = \begin{pmatrix} 0 & \frac{1}{\lambda_k} \\ -m_k \omega_k & 0 \end{pmatrix}$ ,  $C_k = \begin{pmatrix} 0 & 0 \\ -M \omega^2 & 0 \end{pmatrix}$  with boundary conditions  $P\bar{X}(x_0) + Q\bar{X}(x_n) = \bar{0}$ . Cauchy matrix has the following structure

$$\tilde{B}(x, x_0, \omega) \stackrel{def}{=} \sum_{i=0}^{n-1} \tilde{B}_i(x, x_i, \omega) \cdot \tilde{B}(x_i, x_0, \omega) \cdot \theta_i,$$

where  $\tilde{B}(x_i, x_0, \omega) \stackrel{def}{=} \prod_{j=0}^i \tilde{C}_j \cdot \tilde{B}_{i-j}(x_{i-j+1}, x_{i-j}, \omega)$ ,  $\tilde{C}_i = (E + C_i)$ ,  $i = \overline{1, n-1}$ ,

$\tilde{B}(x_i, x_i, \omega) \stackrel{def}{=} E$ ,

$$\tilde{B}_i(x, s, \omega) = \begin{pmatrix} \cos \alpha_i(x - s) & \frac{\sin \alpha_i(x - s)}{\lambda_i \alpha_i} \\ -\lambda_i \alpha_i \sin \alpha_i(x - s) & \cos \alpha_i(x - s) \end{pmatrix}, \alpha_i = \omega \sqrt{\frac{m_i}{\lambda_i}}.$$

Vector  $\bar{C}$  in (14) is  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ .

The characteristic equation of function  $v(x, t)$  takes the form  $b_{12}(\omega) = 0$ . The function  $v(x, t)$  is the same as in (17).

### 7 A Case of Piecewise Constant Functions

Let in (1)  $m(x)$ ,  $f(x)$ ,  $\lambda(x)$  – are piecewise constant functions:

$$m(x) = \sum_{i=0}^{n-1} m_i \theta_i(x), f(x) = \sum_{i=0}^{n-1} f_i \theta_i(x), \lambda(x) = \sum_{i=0}^{n-1} \lambda_i \theta_i(x).$$

Let's consider boundary conditions (6) with matrix  $P = \begin{pmatrix} 1 & 1 \\ 0 & 0 \end{pmatrix}$ ,  $Q = \begin{pmatrix} 0 & 0 \\ 1 & 1 \end{pmatrix}$ .

Under these conditions, the Cauchy matrix components  $b_i(x, s) = \frac{x-s}{\lambda_i}$ .

Let's denote

$$\begin{aligned} \sigma_n &= q_{12} \sum_{m=0}^{n-1} b_m(x_{m+1}, x_m) + q_{22}, \\ I_{k-1}(x_k) &= - \int_{x_{k-1}}^{x_k} b_{k-1}(x_k, s) f_{k-1} ds, \\ I_{k-1}^{[1]}(x_k) &= - \int_{x_{k-1}}^{x_k} f_{k-1} ds. \end{aligned}$$

The function  $v(x, t)$  is the following

$$\begin{aligned} v(x, t) &= \frac{1}{p_{11}\sigma_n - q_{21}p_{12}} \cdot (\psi_0(t)\sigma_n - p_{12}\psi_n(t) - \psi_0(t)q_{21} \cdot (b_i(x, x_i) \\ &+ \sum_{m=0}^{i-1} b_m(x_{m+1}, x_m))) + \frac{p_{12}}{p_{11}\sigma_n - q_{21}p_{12}} \cdot (\psi_n(t) - q_{21}(\sum_{k=0}^n I_{k-1}(x_k) \\ &+ I_{k-1}^{[1]}(x_k) \cdot \sum_{m=k}^{n-1} b_m(x_{m+1}, x_m)) - q_{22} \sum_{k=0}^n I_{k-1}^{[k]}(x_k)) \cdot \\ &(1 + b_i(x, x_i) + \sum_{m=0}^{i-1} b_m(x_{m+1}, x_m)) \end{aligned}$$

In this case the characteristic equation of the problem to eigenvalues is

$$b_{12}(\omega) + b_{22}(\omega) - b_{11}(\omega) - b_{21}(\omega) = 0.$$

Vector  $\bar{C}$  in (14) is  $\begin{pmatrix} -1 \\ 1 \end{pmatrix}$ . The function  $v(x, t)$  is the same as in (17).

By specifying the number of partition segments, material parameters, and core dimensions, we can obtain an analytical expression of the required number of eigenvalues and eigenfunctions. We do all the calculations in Maple package.

### 8 An Example of a Numerical Implementation of the Method for a Rod of Two Pieces

Modern software allows you to get the required number of eigenvalues and eigenfunctions, which ensures the appropriate accuracy of the solution. Consider the result of using the Maple package to obtain a solution to the problem. For example, consider a steel rod 1 m long, consisting of two cylindrical pieces of equal length cross-sectional area, respectively, are  $F_0 = 0,0025 \pi m^2$ ,  $F_1 = 0,000625 \pi m^2$ ,  $x_0 = 0$ ,  $x_1 = 0,5$ ,  $x_2 = 1$ . The Young's modulus for steel is

$E = 20394324259 \text{ kg/m}^2$ , density is  $\rho = 7900 \text{ kg/m}^3$ . Consider the equation of longitudinal oscillations of the rod

$$\frac{\rho}{E} \cdot F(x) \frac{\partial^2 u}{\partial t^2} = \frac{\partial}{\partial x} \left( F(x) \frac{\partial u}{\partial x} \right)$$

with boundary conditions

$$\begin{cases} u(x_0, t) = 1, \\ u(x_2, t) = 1 \end{cases}$$

and initial conditions

$$\begin{cases} u(x, 0) = \varphi_0(x), \\ \frac{\partial u}{\partial t}(x, 0) = \varphi_1(x). \end{cases}$$

Calculations are performed in the Maple package (Figs. 1, 2, 3, 4 and 5).

```

Finding the first eleven eigenvalues  $\omega_k$ 
>  $\omega_1 := \text{NextZero}(\omega \rightarrow b_{12}(\omega), 0, \text{guardDigits} = 500, \text{maxdistance} = 1000000000000)$ ; for k from 1 to 10 do  $\omega_{k+1} := \text{NextZero}(\omega \rightarrow b_{12}(\omega),$ 
 $\omega_k, \text{guardDigits} = 500, \text{maxdistance} = 1000000000000)$  od
 $\omega_1 := 5047.6704$ 
 $\omega_2 := 10095.341$ 
 $\omega_3 := 15143.011$ 
 $\omega_4 := 20190.682$ 
 $\omega_5 := 25238.352$ 
 $\omega_6 := 30286.022$ 
 $\omega_7 := 35333.693$ 
 $\omega_8 := 40381.363$ 
 $\omega_9 := 45429.033$ 
 $\omega_{10} := 50476.704$ 
 $\omega_{11} := 55524.374$ 

```

**Fig. 1.** Finding the first eleven eigenvalues

```

Finding the first eleven eigenfunctions  $X_{k,0}$ 
>  $\text{evalm}\left(\begin{bmatrix} 1 & 0 \end{bmatrix} \& * \text{Matrix}\left(\left[\left[\cos(\alpha \cdot (x-s)), \frac{\sin(\alpha \cdot (x-s))}{\alpha \cdot F_0}\right], [-\alpha \cdot F_0 \cdot \sin(\alpha \cdot (x-s)), \cos(\alpha \cdot (x-s))]\right]\right) \& * \begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) : Y_{k0} := \text{evalf}(\text{subs}(s$ 
 $= x_0, \%))$ ;
 $Y_{k0} := \frac{\sin(\alpha(x-1, x_0))}{\alpha F_0}$ 
> for k from 1 to 11 do  $X_{k,0} := \text{subs}(\alpha = \alpha_k, Y_{k0})$  od
 $X_{1,0} := 40.528473 \sin(3.1415926 x)$ 
 $X_{2,0} := 20.264236 \sin(6.2831855 x)$ 
 $X_{3,0} := 13.509491 \sin(9.4247778 x)$ 
 $X_{4,0} := 10.132118 \sin(12.566371 x)$ 
 $X_{5,0} := 8.1056945 \sin(15.707963 x)$ 
 $X_{6,0} := 6.7547453 \sin(18.849556 x)$ 
 $X_{7,0} := 5.7897816 \sin(21.991149 x)$ 
 $X_{8,0} := 5.0660590 \sin(25.132741 x)$ 
 $X_{9,0} := 4.5031637 \sin(28.274333 x)$ 
 $X_{10,0} := 4.0528473 \sin(31.415926 x)$ 
 $X_{11,0} := 3.6844066 \sin(34.557519 x)$ 

```

**Fig. 2.** Finding the first eleven eigenfunctions  $X_{k,0}$

```

Finding the first eleven eigenfunctions  $X_{k,1}$ 
> evalm( [ 1 0 ] &* Matrix( [ [ cos(α·(x-s)), sin(α·(x-s)) ], [ -α·F1·sin(α·(x-s)), cos(α·(x-s)) ] ] ) &* B &* [ 0 ] ] : Yk,l
:= evalf( subs( s = x1 % ) );
Yk,l :=  $\frac{1.0228719 \cdot 10^6 \cos(\alpha(x-0.5)) \cos(0.00031119234 \sqrt{\omega^2}) \sin(0.00031119234 \sqrt{\omega^2})}{\sqrt{\omega^2}}$ 
+  $\frac{509.29581 \sin(\alpha(x-0.5)) (-0.24999999 \sin(0.00031119234 \sqrt{\omega^2})^2 + \cos(0.00031119234 \sqrt{\omega^2})^2)}{\alpha}$ 
> for k from 1 to 11 do Xk,1 := evalf( subs( α = αk, ω = ωk, Yk,l ) ) od
X1,1 := -0.000014834456 cos(3.1415926 x - 1.5707963) - 40.528471 sin(3.1415926 x - 1.5707963)
X2,1 := 0.000014834455 cos(6.2831855 x - 3.1415928) + 81.056944 sin(6.2831855 x - 3.1415928)
X3,1 := -0.0000013249643 cos(9.4247778 x - 4.7123889) - 13.509491 sin(9.4247778 x - 4.7123889)
X4,1 := 0.000014834455 cos(12.566371 x - 6.2831855) + 40.528472 sin(12.566371 x - 6.2831855)
X5,1 := -0.0000067287610 cos(15.707963 x - 7.8539815) - 8.1056944 sin(15.707963 x - 7.8539815)
X6,1 := 0.0000047023372 cos(18.849556 x - 9.4247780) + 27.018982 sin(18.849556 x - 9.4247780)
X7,1 := -0.000020624237 cos(21.991149 x - 10.995574) - 5.7897815 sin(21.991149 x - 10.995574)
X8,1 := 0.0000097683964 cos(25.132741 x - 12.566370) + 20.264237 sin(25.132741 x - 12.566370)
X9,1 := -0.0000013249643 cos(28.274333 x - 14.137166) - 4.5031636 sin(28.274333 x - 14.137166)
X10,1 := 0.000014834456 cos(31.415926 x - 15.707963) + 16.211389 sin(31.415926 x - 15.707963)
X11,1 := -0.0000074656420 cos(34.557519 x - 17.278760) - 3.6844066 sin(34.557519 x - 17.278760)
    
```

Fig. 3. Finding the first eleven eigenfunctions  $X_{k,1}$




Fig. 4. Finding the first eleven eigenfunctions  $X_{k,0}$




Fig. 5. Finding the first eleven eigenfunctions  $X_{k,1}$

## 9 Conclusions

In the work on the basis of the created new method of solving nonstationary problems for equations of hyperbolic type, the solution of the actual scientific and technical problem by methods of mathematical modeling of wave processes is given. The method of solving equations in partial derivatives of the second order of the hyperbolic type described in the work makes it possible to model oscillating processes in horizontal rods consisting of an arbitrary number of pieces and having different cross sections. The proposed direct method can be used in the study of oscillatory processes without the use of approximate and operational calculus methods. The solutions of the hyperbolic equation with piecewise continuous coefficients on the spatial variable and right-hand sides with the most general local boundary conditions are obtained. The partial case, piecewise-constant coefficients and right-hand sides are singled out, when the solutions of the initial problem can be obtained in a closed form. Using the reduction method, the solutions

of the general first boundary value problem for the equation of hyperbolic type with piecewise continuous coefficients and stationary inhomogeneity are obtained. The general first boundary value problem for the equation of hyperbolic type with piecewise - constant coefficients and - features is investigated. By specifying the number of partition segments, material parameters, and core dimensions, we can obtain an analytical expression of the required number of eigenvalues and eigenfunctions. We do all the calculations in Maple package. The possibilities of application of the proposed method are much wider than this work and, in particular, can be used in further research.

## References

1. Al-Khaled, K., Hazaimah, H.: Comparison methods for solving non-linear sturm-liouville eigenvalues problem. *Symmetry* **1179**(12), 1–17 (2020). <https://doi.org/10.3390/sym12071179>
2. Arsenin, V.Y.: *Methods of Mathematical Physics and Special Functions*, p. 432. Nauka, Moscow (1984)
3. Ashyralyev, A., Aggez, N.: Nonlocal boundary value hyperbolic problems involving integral conditions. *Bound. Value Prob.* (1), 1–10 (2014). <https://doi.org/10.1186/s13661-014-0205-4>
4. Atkinson, F.: *Discrete and Continuous Boundary Value Problems*, p. 518. Academic Press, Cambridge (1964)
5. Borwein, J.M., Skerritt, M.P.: *An Introduction to Modern Mathematical Computing: With Maple*, p. 233. Springer, New York (2011). <https://doi.org/10.1007/978-1-4614-0122-3>
6. Hornikx, M.: The extended fourier pseudospectral time-domain method for atmospheric sound propagation. *J. Acoust. Soc. Am.* **1632**(4), 1–20 (2010). <https://doi.org/10.1121/1.3474234>
7. Kong, O., Wu, H., Zettl, A.: Sturm-liouville problems with finite spectrum. *J. Math. Anal. Appl.* **263**, 748–762 (2001). <https://doi.org/10.1006/jmaa.2001.7661>
8. Lysenko, A., Yurkov, N., Trusov, V., Zhashkova, T., Lavendels, J.: Sum-of-squares based cluster validity index and significance analysis. *Lect. Notes Comput. Sci.* (including subseries *Lecture Notes in Artificial Intelligence and Lecture Notes in Bioinformatics*) **5495**, 313–322 (2009). [https://doi.org/10.1007/978-3-642-04921-7\\_32](https://doi.org/10.1007/978-3-642-04921-7_32)
9. Martin, N., Nilsson, P.: The moving-eigenvalue method: hitting time for ito processes and moving boundaries. *J. Phys. A Math. Theoretica* **53**(40), 1–32 (2020). <https://doi.org/10.1088/1751-8121/ab9c59>
10. Mennicken, R., Möller, M.: *Non-Self-Adjoint Boundary Eigenvalue Problems*, p. 518. North Holland (2003)
11. Mukhtarov, O., Yücel, M.: A study of the eigenfunctions of the singular sturm-liouville problem using the analytical method and the decomposition technique. *Mathematics* **415**(8), 1–14 (2020). <https://doi.org/10.3390/math8030415>
12. Sabitov, K.B., Zaitseva, N.V.: Initial-boundary value problem for hyperbolic equation with singular coefficient and integral condition of second kind. *Lobachevskii J. Math.* **39**(9), 1419–1427 (2018). <https://doi.org/10.1134/S1995080218090299>
13. Tatsii, R.M., Pazen, O.Y.: Direct (classical) method of calculation of the temperature field in a hollow multilayer cylinder. *J. Eng. Phys. Thermophys.* **91**(6), 1373–1384 (2018). <https://doi.org/10.1007/s10891-018-1871-3>



14. Tatsij, R.M., Chmyr, O.Y., Karabyn, O.O.: The total first boundary value problem for equation of hyperbolic type with pieewise constant coefficients and delta-singularities. *Res. Math. Mech.* **24**, 86–102 (2019)
15. Tichonov, A., Samarskii, A.: *Equations of Mathematical Physics*, chap. 2: Equations of the Hyperbolic Type, p. 777. Pergamon Press, Oxford (1990)
16. Tisseur, F., Meerbergen, K.: The quadratic eigenvalue problem. *SIAM Rev.* **42**(2), 235–286 (2001). <https://doi.org/10.1137/S0036144500381988>
17. Wyld, H.W., Powell, G.: *Mathematical Methods for pPhysics*, Chap. 1: Homogeneous Boundary Value Problems and Special Functions, p. 476. CRC Press, Boca Raton (2020)
18. Yang, F., Zhang, Y., Liu, X., Li, X.: The quasi-boundary value method for identifying the initial value of the space-time fractional diffusion equation. *Acta Mathematica Scientia* **40**(3), 641–658 (2020). <https://doi.org/10.1007/s10473-020-0304-5>
19. Yarka, U., Fedushko, S., Veselý, P.: The dirichlet problem for the perturbed elliptic equation. *Mathematics* **8**, 1–13 (2020). <https://doi.org/10.3390/math8122108>